

Development Team

Principal Investigator Dr. Sunil Kumar Khare, Professor, Department of Chemistry, IIT-Delhi

Content Writer: Dr. Archana Sachdeva, Professor, Division of Biochemistry, Indian Agricultural Research Institute (IARI), New Delhi

Content Reviewer: Dr. Tanmay Dutta, Assistant Professor, Department of Chemistry, Indian Institute of Technology, Delhi

Paper Coordinator: Dr. Archana Sachdeva, Professor, Division of Biochemistry, Indian Agricultural Research Institute (IARI), New Delhi

Description of Module

Subject Name	Biochemistry
Paper Name	08 Protein Biochemistry and Enzymology
Module Name/Title	Enzyme Inhibition

 **Pathshala**
पाठशाला
A Gateway to All Post Graduate Courses

Biochemistry

Protein Biochemistry and enzymology

Enzyme Inhibition

1. Objectives

- Enzyme inhibition
- Various types of enzyme inhibitions

2. Description – Enzyme Inhibition

An enzyme catalyzed reaction can be hindered or reduced by a number of substances. Some others like urea are known as denaturants, being non-specific in their mode of action. But if any compound act in a fairly specific way in inhibiting the catalysis of a particular enzyme they are called **inhibitors**. The loss in activity can either be of two types-

- (1) Reversible, where the activity can be restored by the removal of the inhibiting compound. It's temporary.
- (2) Irreversible, where the loss of activity cannot be recovered within the stipulated time of interest. It is permanent. Irreversible inhibition behave as time dependent loss of enzyme concentration with lowered V_{max} or incomplete in activation with time dependent change in both K_m and V_{max} .

Heavy metal ions like mercury, lead etc cause irreversible inhibition, which bind strongly to the amino acid backbone termed as "**suicide inhibition**".

There are several reasons behind the need for studying enzyme inhibition mechanisms. They are:-

- ❖ Exploring potential mechanisms in multi-substrate reactions.
- ❖ Studying the relative binding affinity of competitive inhibitors to the enzyme active site, in the absence of 3-D structure information.
- ❖ For understanding various control mechanisms-how the balance of protease enzymes and their inhibitors in tissues achieve homeostasis.
- ❖ For various commercial applications like pesticide, insecticide, weed-killers, pharmaceutical compounds like drugs etc.




Figure 1. Types of enzyme inhibition

2.1. Reversible Inhibition:

In this type of inhibition, the hindrance is temporary and thus noncovalent interactions like hydrogen bonds, ionic bonds or hydrophobic bonds form between inhibitors and the enzyme. Even though these are weak bonds, multiple such bonds cause strong and specific binding. Despite absence of any chemical reactions, the inhibitors can easily remove or exchange by dilution or dialysis. After removing the inhibitor, enzyme can be fully restored in reversible inhibition. Equilibrium is established between free inhibitor and enzyme-inhibitor [EI] complex (Figure 2).




Figure 2. Mechanism of reversible inhibition

Reversible inhibitions are of different types. The classification is based according to the effect of varying the concentration of the enzyme's substrate on the inhibitor.

2.1.1. **Competitive Inhibition:** In this type of reversible inhibition, both the substrate and its inhibitor cannot bind to the enzyme at the same time to the allosteric/active site. This normally occurs due to the structural similarity of substrate and the inhibitor, which results with affinity for the active site. The inhibition can be recovered by the presence of high concentration of substrate, outcoming the competing inhibitor. V_{max} of the reaction is unchanged, while K_d , the dissociation constant is apparently increased. Competitive inhibitors can also be used to find the enzyme active site.

Eg: N-(phosphonacetyl)-L-aspartate also known as PALA is a competitive inhibitor for aspartate transcarbamoylase.

Eg: Malonate is a competitive inhibitor of enzyme succinate dehydrogenase, and competes with succinate.




Figure 3. Kinetics of competitive inhibition

Table 1. Clinical use of competitive inhibition

Drugs	Target Enzyme	Therapeutic use
STATINS , Atorvastatin Simvastatin	HMG CoA reductase	Involved in the reduction of plasma cholesterol level- Anti-hyperlipidemic agents
Allopurinol	Xanthine oxidase	Used in case of prevention

		of gout attacks
Methotrexate	Dihydrofolate reductase	Used as cancer treatment drug
Captopril, Enalapril	Angiotensin converting enzyme	Treatment of high blood pressure
Discoumarol	Vit K-epoxide reductase	Used as an anti-coagulant

2.1.2. **Uncompetitive Inhibition:** It's an anti-competitive inhibition; where the inhibitor binds only to the substrate-enzyme complex. According to its kinetics, V_{max} and K_m decrease. This type of inhibition works best in case of high concentration of the substrate. The substrate and the uncompetitive inhibitor does not resemble each other. Eg:-Lithium and phosphoinositide cycle.




Figure 4. Kinetics of un-competitive inhibition

2.1.3. **Non-competitive Inhibition:** A non-competitive inhibitor is one which reacts with enzyme-substrate or [ES] complex. It does not affect the binding of the substrate, but slows down the reaction rate for formation of the enzyme-product [EP] complex. The only factor on which the extent of hindrance or inhibition depends is the inhibitor concentration. There will be a decrease in V_{max} but k_m will remain the same.

- Eg: Alanine non competitively inhibits the enzyme pyruvate kinase.




Figure 5. Kinetics of non-competitive inhibition

Table 2. Examples of common non-competitive inhibitors

Inhibitor	Enzyme inhibited
Heavy metals – Ag^{2+} , Hg^{2+} , Pb^{2+}	Heavy metals bind with cysteinyl SH group of enzyme
Pepstatin	Pepsin
Soybean trypsin inhibitor	Trypsin
Ethanol/narcotic drugs	Acid phosphatase

2.1.4. **Mixed Inhibition:** In this type of inhibitor, inhibitor is capable of binding to both free enzyme as well as enzyme- substrate complex. In this case, V_{\max} and k_{\max} varies. Mixed inhibitor binds to the allosteric site. This type of inhibition cannot overcome by increasing substrate concentration S , but can be reduced. The inhibitor binding to the allosteric site changes the structural confirmation to reduce the affinity of the substrate. Eg: Mixed inhibition is observed on case of oxidoreductase activity of xanthine.

oxidase by Pd²⁺ ion

$$\frac{1}{v} = \frac{k_s}{v_{\max}} \left(1 + \frac{[I]}{k_i}\right) \frac{1}{[S]} + \frac{1}{v_{\max}} \left(1 + \frac{[I]}{\alpha k_i}\right)$$




Figure 6. Kinetics of mixed inhibition




Figure 7. Different types of reversible inhibitions

2.2. Irreversible Inhibition: In this type of inhibition, the hindrance is of permanent nature by modifying enzyme covalently. These types of inhibitors often contain electrophilic functional groups like fluorophosphates, aldehydes, haloalkanes, alkenes, nitrogen mustards, phenyl sulfonates, Michael acceptors etc, which react with aminoacid sidechains having nucleophilic residues.

These inhibitors are very specific in the mechanism of inactivation for a particular class of enzyme-They do irreversible inhibition by specially altering the active site. They display inhibition which is time-dependent. Their potency cannot be characterized by IC_{50} value. These inhibitors increase K_m and decrease V_{max} .

Eg: Diisopropylfluorophosphate (DFP) is an irreversible protease inhibitor.




Figure 8. Kinetics of irreversible inhibition

Table 3. Therapeutic uses of irreversible inhibitors.

Inhibitors	Enzyme inhibited	Therapeutic uses
Disulfiram	Aldehyde dehydrogenase	Treats alcoholism
Cyanide ions	Cytochrome oxidase	Inhibition of the respiratory chain

Fluoride ions	Enolase	Inhibition of Glycolysis
Melathion	Acetyl choline esterase	Used as an organophosphorus insecticide
Di-isopropyl fluorophosphate	Serine proteases, Acetyl choline esterase	Used as a nerve gas
British Anti Lewisite (BAL)	Reaction with the thiol (-SH) group of the enzyme	Used as an antidote in case of poisoning due to heavy metal

Suicide inhibition- This is another type of irreversible inhibition. In this case, the target enzyme converts the inhibitor compound into a reactive form in its active site. They are also known as mechanism based inhibitors or transition state analogs.

Eg:- DFMO [α -difluoromethyl ornithine], an analogue of ornithine inhibits ornithine decarboxylase.

Eg:- Allopurinol is a suicide inhibitor of xanthine oxidase Aspirin inhibits cyclooxygenase.

Table 4. Therapeutic uses of suicide inhibitors

Drugs	Product	Target Enzyme	Therapeutic use
Allopurinol	Alloxanthin	Xanthine oxidase	Treatment of gout
5-Fluorouracil	Fluoro-deoxy uridylate	Thymidylate synthase	Cancer treatment
Aspirin	Acetylates the serine residue present in the cyclooxygenase active site	Cyclooxygenase	Used as a non-steroidal anti-inflammatory drug
Difluoro methyl ornithine (DFMO)	Forms an irreversible covalent complex with the co-enzyme	Ornithine decarboxylase	Used for treating Sleeping sickness (trypanosomiasis)

Table 5. Differences between reversible & irreversible inhibitions

Reversible	Irreversible
Binds via non-covalent interactions	Binds via covalent interactions
Do not perform any chemical changes	Inhibitor binds to the substrate and prevent catalytic activity of enzymes
As there is no bonding between the inhibitor and substrate, reversible inhibition can be reversed,	Irreversibility due to strong covalent bonding

2.3. **Allosteric Inhibition:** Allosteric inhibition is a type of enzyme regulation, in which allosteric inhibitor binds to a site other than the active site of the enzyme. This additional site to which effector binds is called **allosteric site**. When these effectors bind to the protein, results with conformational change and cause enhancement in activity is known as allosteric activators. When they decrease the activity of the protein, they are known as **allosteric inhibitors**. Allosteric enzymes are K or V types.

Models of allosteric regulation: The allosteric effects or mechanism is well described by the concerted HWC model, which was put forth by Monod, Wyman and Changeux. Another model called the sequential model, proposed by Koshland, Nemethy and Filmer, also possibly explains the allosteric regulation. Both these models postulate that enzyme subunits exist in one two conformations - tensed (T) or relaxed(R) states.

2.3.1 Concerted model: This model is known as symmetry or MWC model. According to this model, enzyme subunits exist in same conformation, they are connected and a slight conformational change in any one of the subunits is conferred to all other subunits of the enzyme. When any ligand or substrate is absent, the equilibrium favors towards either of the conformational states. Among the tensed and relaxed states, the 'R' state has higher

affinity than 'T' state. The most successful application of this model is this regulation of hemoglobin.

2.3.2. Sequential model: In contrary to concert model, sequential model states that enzymes subunits are not connected, such that any change in the enzyme conformation leads to induction of a similar change in the others. When a subunit randomly collides with substrate, an induced fit converts a subunit from the 'T' state to 'R' state.

2.3.2. Morphein model: The third model is a dissociative concerted model known as morphein model. This is physiologically significant homo-oligomeric tetramer structure. Transition in the morphein model are assisted by dissociation of oligomer, conformational change in dissociated state and reassembly of oligomers. So far, one of the best characterized morphein is the enzyme porphobilinogen synthase.

Types of allosteric regulation:

There are mainly two types of allosteric regulation. Homotropic & Heterotropic.




Figure 9. Types of allosteric regulation

- **Homotropic regulation:** It's a positive modulation- The modulator acts not only as a substrate but also as a regulatory molecule of the target enzyme.
Eg: CO₂ is a heterotropic modulator of haemoglobin.
- **Heterotropic regulation**It can either be a positive or negative modulation. Here the modulator is a regulatory molecule but not an enzymes substrate.
Eg:CO₂is a heterotropic modulator of haemoglobin.

3. Importance of enzyme inhibition:

- ✓ Understanding regulation of enzyme activity in living cells
- ✓ Elucidation of the cellular metabolic pathways by accumulation of intermediates
- ✓ Helps in identification of catalytic or functional groups present at the enzyme active site
- ✓ Helps in providing information on enzyme's substrate specificity
- ✓ Helps in studying the mechanism of catalytic activity
- ✓ Competitive or suicide inhibitors also find therapeutic applications.

Conclusion

Specific inhibition of target enzyme has been attributed through various modes of inhibition. It can either be reversible, irreversible or allosteric model of inhibitions. Other than the models and mechanisms of inhibitions, they have immense applications.